

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 098 098
A2

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 83303610.6

(22) Date of filing: 22.06.83

(51) Int. Cl.³: **C 07 D 219/04**
C 07 D 219/10, C 07 D 219/06
A 61 K 31/435

(30) Priority: 25.06.82 NZ 201084

(43) Date of publication of application:
11.01.84 Bulletin 84/2

(84) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

(71) Applicant: **DEVELOPMENT FINANCE CORPORATION OF NEW ZEALAND**
Development Finance Centre Corner Grey and Featherston Streets
Wellington(NZ)

(72) Inventor: **Atwell, Graham John**
37 Hawkins Street Meadowbank
Auckland 5(NZ)

(72) Inventor: **Baguley, Bruce Charles**
36a Olsen Avenue Hillsborough
Auckland 4(NZ)


(72) Inventor: **Denny, William Alexander**
165 Gossamer Drive Pakuranga
Auckland(NZ)

(72) Inventor: **Rewcastle, Gordon William**
24 Rothery Road Manurewa
Auckland(NZ)

(74) Representative: **Senior, Janet et al,**
Abel & Imray Northumberland House 303-306 High Holborn
London WC1V 7LH(GB)

(54) Acridinecarboxamide compounds.

(57) The novel class of 4-carboxamidoacridines of the present invention represented by the general formula (I),



where R_1 represents H, CH_3 or NHR_3 , where R_3 is H, $COCH_3$, SO_2CH_3 , $COPh$, SO_2Ph or lower alkyl optionally substituted with hydroxyl and/or amino functions;

R_2 represents H or up to two of the groups CH_3 , OCH_3 , halogen, CF_3 , NO_2 , NH_2 , $NHCOCH_3$, and $NHCOOCH_3$ placed at positions 1-3 and 5-8;


Y represents $.C(NH)NH_2$, $NHC(NH)NH_2$, or NR_4R_5 , where each of R_4 and R_5 is H or lower alkyl optionally substituted with hydroxyl and/or amino functions; and

x is from 2 to 6,

and the acid addition salts thereof, possess antibacterial and antitumour properties.

The present invention relates to novel acridine derivatives having antibacterial and antitumour properties, to methods for preparing these compounds, and to the use of the compounds as antibacterial and antitumour agents. The present invention also relates to novel compounds useful as intermediates in the preparation of the acridine derivatives of the invention.

The novel class of 4-carboxamidoacridines of the present invention is represented by the general formula (I),



where R_1 represents H, CH_3 or NHR_3 , where R_3 is H, $COCH_3$, SO_2CH_3 , $COPh$, SO_2Ph or lower alkyl optionally substituted with hydroxyl and/or amino functions;

R_2 represents H or up to two of the groups CH_3 , OCH_3 , halogen, CF_3 , NO_2 , NH_2 , $NHCOCH_3$, and $NHCOOCH_3$ placed at positions 1-3 and 5-8;

Y represents $C(NH)NH_2$, $NHC(NH)NH_2$, or NR_4R_5 , where each of R_4 and R_5 is H or lower alkyl optionally substituted with hydroxyl and/or amino functions; and

x is from 2 to 6,

and the acid addition salts thereof.

When R_3 , R_4 or R_5 represent lower alkyl, the group may contain from 1 to 4 carbon atoms.

A preferred subclass of these compounds of formula (I) are those where R_1 represents NH_2 , R_2 represents up to two of 1-, 5-, 6-, 7- or 8- NO_2 , 5- or 6- CH_3 , and 5- Cl , Y represents $NHC(NH)NH_2$, $N(CH_3)_2$, or $NHCH_2CH_2OH$ and x is 2.

Another preferred subclass of these compounds of formula (I) has the same values for R_2 , Y and x but R_1 represents H .


Four specific compounds of formula (I) are those in which,

- (a) R_1 and R_2 represent H , Y represents $N(CH_3)_2$ and x is 2;
- 5 (b) R_1 represents NH_2 , R_2 represents H , Y represents $N(CH_3)_2$ and x is 2;
- (c) R_1 represents NH_2 , R_2 represents $6-NO_2$, Y represents $N(CH_3)_2$ and x is 2; and
- 10 (d) R_1 represents NH_2 , R_2 represents $5-CH_3$, Y represents $N(CH_3)_2$ and x is 2.

Other specific compounds of formula (I) are listed in Tables I and II hereinafter.

15 The compounds of formula (I) form pharmaceutically acceptable addition salts with both organic and inorganic acids. Examples of suitable acids for salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic, and the like.

20 The compounds of general formula (I) and the acid addition salts thereof may be prepared for example by a process which comprises coupling a substituted acridine of the general formula (II),



II

where R_2 is as defined as above, Z represents H , CH_3 , or any suitable leaving group (e.g. methoxy, phenoxy, alkylthio or halogen, but preferably chloro) and R_6 represents Cl , Br or

OC₆H₄-p-NO₂, with a primary alkyl amine of the general formula (III),

III

where x and Y are as defined above, and, when Z is a leaving group, converting the resultant coupled product to a compound of general formula (I) where R₁ represents NHR₃ and R₃ is as defined above, and, if desired, converting a compound of formula (I) into an acid addition salt thereof.

The coupling reaction is desirably performed in an anhydrous solvent (e.g. chloroform, dimethylsulphoxide or N-methylpyrrolidone, but preferably dichloromethane or dimethylformamide) buffered with a tertiary amine, preferably triethylamine. The reaction is conveniently performed at temperatures in the range from 0°C to 50°C, with the preferred temperature being 20°C.

In the case of Z representing Cl in formula (II), further treatment of the resulting coupled products with anhydrous ammonia or suitable amine of the general formula R₃NH₂ in phenol or cresol provides the compounds of formula (I) where R₁ represents NHR₃. Alternatively, treating the resulting coupled products where Z=Cl with neat phenol or cresol provides corresponding compounds where Z=OC₆H₅ or OC₆H₄CH₃ and R₆ is NH(CH₂)_xY, where x and Y are defined as for formula (I). These compounds can be isolated, but are usually treated in situ with anhydrous ammonia or amine R₃NH₂ to provide the desired compounds of general formula (I).


The acid addition salts of the compounds of formula (I) may be prepared for example by contacting the free base form with an equivalent amount

of the desired acid in the conventional manner. The free base forms may be regenerated by treating the salt form with a base. For example, dilute aqueous base solutions may be utilized. Dilute
5 aqueous potassium hydroxide, potassium carbonate, ammonia and sodium bicarbonate solutions are for example suitable for this purpose. The free base forms differ from their respective salt forms somewhat in certain physical properties such as solubility
10 in polar solvents, but in general the salts are otherwise equivalent to their respective free base forms for the purposes of the invention.

The primary alkyl amines of the general formula (III) are known compounds and are commercially available or preparable by methods
15 described in the literature. Examples of such compounds include N,N-dimethyl-1,2-ethanediamine (N,N-dimethylethylenediamine), N,N-diethyl-1,2-ethanediamine, N,N-dimethyl-1,3-propanediamine, N,N-dimethyl-1,4-butanediamine, N,N-dimethyl-1,5-pentanediamine, N-(2-hydroxyethyl)-1,2-ethanediamine
20 (2-(2-aminoethylamino)-ethanol), N-methyl-N-(2-hydroxyethyl)-1,2-ethanediamine, 2-aminoethylguanidine $\text{NH}_2(\text{CH}_2)_2\text{NHC}(\text{NH})\text{NH}_2$, and 3-aminopropionamidide $\text{NH}_2(\text{CH}_2)_2\text{C}(\text{NH})\text{NH}_2$. The two last-mentioned compounds may be prepared according to P.L. Barker, P.L. Gendler,
25 and H. Rapoport, J.Org.Chem., 46, 2455 (1981).

The amines of the general formula R_3NH_2 are also known compounds, and are commercially available or preparable by methods described in the literature. Examples of such compounds where R_3 is lower alkyl
30 optionally substituted with hydroxyl and/or amino functions include methylamine, ethylamine, 2-hydroxyethylamine, 2,3-dihydroxypropylamine, and N,N-dimethyl-1,2-ethanediamine (N,N-dimethylethylenediamine).


The 9-substituted acridines of formula (II) are novel compounds useful as intermediates in the preparation of the compounds of formula (I), and accordingly the present invention also provides the compounds represented by the general formula (II),



II

where R_2 is defined as for formula (I), Z represents H, CH_3 , or any suitable leaving group (e.g. methoxy, phenoxy, alkylthio or halogen but preferably chloro) and R_6 represents Cl, Br or $OC_6H_4-p-NO_2$, and acid addition salts thereof.


- 5 The 9-substituted acridines of general formula (II) where Z represents H or halogen may be prepared for example by the process outlined in Scheme I, and this general process also forms part of the present invention. In Scheme I, R_2 is as defined for formula (I), and R_6 is as defined for formula (II).



- 10 Thus, for example:
The diphenylamine diacids (IV) are formed by the Jourdan-Ullmann reaction between suitably substituted 2-halobenzoic acids and anthranilic acids in high yield (see B.F. Cain, G.J. Atwell and W.A. Denny, J.Med.Chem., 20, 987 (1977) and European Patent Application No. 82304420.1). The resulting diphenylamine diacids (IV) are cyclodehydrated with mineral acids or their derivatives (e.g. H_2SO_4 , polyphosphoric acid or polyphosphate ester) to form
- 15

carboxyacridanones, from which the desired 4-carboxy isomers (V) are obtained if necessary by separation from co-occurring isomers. Such separations can readily be achieved by taking advantage of the differential solubilities of the different isomers, both as free acids and acid salts.


Alternatively, the formation of unwanted isomers in the cyclodehydration reaction can be avoided e.g. by the process outlined in Scheme II, and this general process also forms part of the present invention. In Scheme II, R₂ is as defined for formula (I), and R₇ represents a lower alkyl group, i.e. containing from 1 to 4 carbon atoms, preferably methyl or t-butyl, and X is a halogen, preferably Cl or Br, or phenyl-iodonium.



Thus, for example:

The diphenylamine esters (VIII) are formed via a novel modification of the Jourdan-Ullmann reaction, in which a suitable organic base (e.g. tri-*n*-butylamine, N-ethylmorpholine or diisopropylethylamine) is used as both solvent and acid acceptor, thus preventing hydrolysis of the ester group. A soluble form of copper catalyst such as copper (II) acetate is used. Ring closure of the diphenylamine esters (VIII) is effected without concomitant hydrolysis of the ester function by using polyphosphate ester (PPE) as reagent. Subsequent acid- or base-catalysed hydrolysis of the acridone esters (IX) gives the desired products (V).


The acridone esters represented by the general formula (IX),



IX

where R_2 is as defined for formula (I) and R_7 represents a lower alkyl group, preferably methyl or t-butyl, are novel compounds useful as intermediates in the preparation of the compounds of formula (I), and they and their acid addition salts accordingly form part of the present invention.

An alternative preparation of the 4-carboxy compounds of formula (V) is outlined in Scheme III, and this general process also forms part of the present invention. In Scheme III, R_2 is as defined for formula (I) and X represents halogen but preferably iodo.



Scheme III

Thus, for example:

Reaction of the halodiacid with the appropriately substituted amine is carried out in an anhydrous solvent such as N-methylpyrrolidone or dimethylsulphoxide (DMSO), but preferably dimethylformamide (DMF), in the presence of acid acceptors such as K_2CO_3 and organic bases (preferably N-ethylmorpholine) and Cu powder to give the N-substituted diacid (XI). These are conveniently isolated by diluting the reaction mixture with water and extracting with suitable organic solvents, preferably ethyl acetate. The resulting diphenylamine diacids are cyclodehydrated

with mineral acids or their derivatives as described above to form carboxyacridanones, from which the desired 4-carboxyacridanones (V) are obtained if necessary by separation from co-occurring isomers as described above.

- 5 Reduction of the substituted 4-carboxyacridanones (V) to substituted 4-carboxyacridines (VI, Z=H), in Scheme I, can be achieved e.g. by direct treatment with Al/Hg amalgam (A. Albert and E. Ritchie, J.Soc.Chem Ind., 60, 120 (1941), or by formation of the tosylhydrazide adduct (VI, Z=NHNHSO₂C₆H₄CH₃) via the
- 10 corresponding 9-chlorocompound, (VI, Z=Cl), and subsequent base-catalyzed decomposition of the adduct (A. Albert and A. Royer, J.Chem.Soc., 1148, (1949)). Reaction of the 4-carboxyacridanones (V) with tris(4-nitrophenyl)phosphite in pyridine gives the 4-nitrophenylester derivatives (VII) (B.F. Cain, G.J. Atwell and
- 15 W.A. Denny, J.Med.Chem., 20, 987 (1977)). Similar reaction of the 4-carboxyacridines (VI, Z=H) with tris (4-nitrophenyl)phosphite in pyridine gives the compounds of general formula (II) where Z is H and R₆ is OC₆H₄-p-NO₂.

- Compounds of general formula (V) can e.g. be activated by reaction
- 20 with a suitable halogen reagent (e.g. PCl₅, POCl₃, but preferably SOCl₂) and a trace of DMF as catalyst to provide compounds of formula (II) where Z is Cl and R₆ is Cl. Similar activation of compounds of general formula (VI) provides compounds of formula (II) where Z is H and R₆ is Cl. Similar activation of compounds
- 25 of general formula (VII) provides compounds of general formula (II) where Z is Cl and R₆ is OC₆H₄-p-NO₂.

- Similar activation of compounds of general formula (V), (VI) or (VII) with POBr₃ or preferably SOBr₂ provides compounds of formula (II) where Z is Br and R₆ is Br, Z is H and R₆ is Br, or Z
- 30 is Br and R₆ is OC₆H₄-p-NO₂.

These compounds of general formula (II) where Z is H, Cl or Br can then be reacted with amines of general formula (III) e.g. in

anhydrous solvents (e.g. CHCl_3 , DMSO or N-methylpyrrolidone, but preferably CH_2Cl_2 or DMF) buffered with a tertiary amine (preferably triethylamine).

5 The method of preparation when an amine of formula (III) is reacted with a compound of formula (II) where R_6 is $\text{OC}_6\text{H}_4\text{-p-NO}_2$ is the preferred method when the sidechain component (III) contains, in addition to the primary amine, other secondary amine or hydroxylated amine functions Y.


10 The compounds of general formula (II) where Z is alkylthio can be prepared for example by the methods cited in E.F. Elslager et al., J.Med.Chem., 14, 782-788 (1971), and the resultant products from the coupling reaction with amines of general formula (III) can for example be converted to compounds of formula (I) where R_1 is NHR_3 by the methods also cited therein.

15 The compounds of general formula (II) where Z is methoxy or phenoxy can be prepared for example by the methods given in Albert, "The Acridines", Second Edition, Edward Arnold Ltd, London (1966).

20 Other compounds of the general formula II where Z is a leaving group other than the ones specifically listed above may be formed by methods known to the man skilled in the art, for example, where appropriate, by the methods described above.

20

The 9-substituted acridines of general formula (II), wherein Z represents CH_3 may be prepared e.g. by the process outlined in Scheme IV, and this general process also forms part of the present invention. In Scheme IV, X represents halogen but preferably bromo, R_2 and R_6 are as defined for formula (II) and R_8 is as defined below.




Thus, for example:

- Reaction of suitably substituted 2-haloacetophenones with anthranilic acid in the presence of 1 mole of acid acceptor (preferably potassium carbonate) and a catalytic amount of copper gives the
- 5 diphenylamine products (XII), where R_8 is OH. Reaction of the ketoacids (XII; $R_8 = OH$) with a suitable lower alcohol (preferably ethanol) using 1 mole of diethyl phosphorocyanidate (DEPC) or other suitable ester-forming reagents and 1 mole of acid acceptor, preferably triethylamine, gives compounds (XII, $R_8 = OCH_2CH_3$).
- 10 Cyclodehydration, for example using 5% H_2SO_4 in refluxing acetic acid, provides compounds (XIII; $R_8 = OCH_2CH_3$), which can be hydrolyzed in dilute ethanolic sodium hydroxide to the acid (XIII; $R_8 = OH$). Activation of these compounds with a suitable halogen reagent (preferably $SOCl_2$ or $SOBr_2$) as detailed above provides
- 15 compounds of general formula (II) where Z is CH_3 and R_6 is Cl or Br. Reaction of the acid (XIII; $R_8 = OH$) with tris(4-nitrophenyl)phosphite in pyridine gives the 4-nitrophenylester derivatives (II'; $R_6 = OC_6H_4-p-NO_2$) (B.F. Cain, G.J. Atwell and W.A. Denny, J.Med.Chem., 20, 987 (1977)).
- 20 The compounds of general formula (II) where Z is CH_3 and R_6 is Cl, Br or $OC_6H_4-p-NO_2$ may then be coupled with suitable primary amines of general formula (III) e.g. in anhydrous solvent (e.g. $CHCl_3$, DMSO or N-methylpyrrolidone, but preferably CH_2Cl_2 or DMF) buffered with a tertiary amine (preferably triethylamine) to provide
- 25 compounds of general formula (I) where R_1 is CH_3 .

An alternative and preferred process for the preparation of compounds of general formula (I) where R_1 is CH_3 and R_2 is defined as for formula (I) is outlined in Scheme V, and this is also a process of the present invention. In Scheme V, X represents

30 halogen but preferably bromo, R_2 and Y are as defined for formula (I) and R_8 is as defined below.



Thus, for example:

Reaction of suitably substituted 2-haloacetophenones with anthra-
nilic acid in the presence of 1 mole of acid acceptor (preferably
potassium carbonate) and a catalytic amount of copper gives the
5 diphenylamine products (XII), where R_8 is OH. These compounds can
be coupled with a primary alkyl amine of general formula (III) in
anhydrous solvents (e.g. N-methylpyrrolidone or N-methyl aceta-
mide but preferably DMF) using 1 mole of diethyl phosphorocya-
nide or other suitable amide-forming reagent and 1 mole of an
10 acid acceptor, preferably anhydrous triethylamine to give com-
pounds (XII), where R_8 is $\text{NH}(\text{CH}_2)_x\text{Y}$. Cyclodehydration of these
compounds, for example using 5% H_2SO_4 in refluxing acetic acid,
gives the required compounds of general formula (I), where R_1 is
 CH_3 .

- 15 In the various processes described above, the com-
pounds of the general formulae II to XIII may, where
appropriate, be used in the form of their acid
addition salts or in the form of their salts with
bases.
- 20 In the formulae shown above, R_2 represents H or one
or two of the same or different substituents selec-
ted from CH_3 , OCH_3 , halogen, CF_3 , NO_2 , NH_2 , NHCOCH_3
and NHCOOCH_3 . It will be appreciated that, although
the formulae show R_2 as a substituent of one ring
25 of the acridine and diphenylamine derivatives of the

general formulae I, II, IV to IX, XI and XII, when R_2 represents one substituent this may occur on either of the rings and when R_2 represents two substituents these may occur on either or both
5 rings. Thus, for a substituted acridine derivative, the substituent(s) may be at any one or two of the free positions on the rings, i.e. positions 1-3 and 5-8; substituent(s), when present, are correspondingly situated on either or both of the phenyl
10 rings of formulae IV, VIII, XI and XII and these compounds may be derived from appropriately substituted starting materials.

The following Tables I and II set out physical data for 24 compounds within the general formula (I), representative of it, and
15 preparable by the processes of the invention. In Table I the following terms and abbreviations are used:-

MP = melting point of the reported acid addition salt in °C.
R_m = a measure of the compound's lipophilic-hydrophilic balance from reversed phase partition chromatography.
20 R_m is linearly related to partition coefficients obtained in the 1-octanol/water system.

TABLE I

0098098

No	R ₁	R ₂	x	Y	Mp	Formula	Rm
1	H	H	2	N(CH ₃) ₂	195-197	C ₁₈ H ₁₉ N ₃ O.2HCl	-0.20
2	CH ₃	H	2	N(CH ₃) ₂	178-180	C ₁₉ H ₂₁ N ₃ O.2HCl	-0.30
3	NHCH ₃	H	2	N(CH ₃) ₂	231-233	C ₁₉ H ₂₂ N ₄ O.2HCl	-1.11
4	NH ₂	H	2	N(CH ₃) ₂	292-293	C ₁₈ H ₂₀ N ₄ O.2HCl $\frac{1}{2}$ H ₂ O	-1.11
5	NH ₂	H	3	N(CH ₃) ₂	290-292	C ₁₉ H ₂₂ N ₄ O.2HCl	-0.93
6	NH ₂	H	2	NH(CH ₂) ₂ OH	292-293	C ₁₈ H ₂₀ N ₄ O.2HCl	-1.06
7	NH ₂	H	2	N(CH ₂ CH ₃) ₂	283-285	C ₂₀ H ₂₄ N ₄ O.2HCl	-0.67
8	NH ₂	H	2	NH ₂	344-345	C ₁₆ H ₁₆ N ₄ O.2HCl	-1.18
9	NH ₂	2-NO ₂	2	N(CH ₃) ₂	>360	C ₁₈ H ₁₉ N ₅ O ₃ .2HCl	
10	NH ₂	2-NH ₂	2	N(CH ₃) ₂	>360	C ₁₈ H ₂₁ N ₅ O.2HCl	
11	NH ₂	5-NO ₂	2	N(CH ₃) ₂	>360	C ₁₈ H ₁₉ N ₅ O ₃ .2HCl	
12	NH ₂	5-NH ₂	2	N(CH ₃) ₂	326-329	C ₁₈ H ₂₁ N ₅ O.2HCl.H ₂ O	
13	NH ₂	5-CH ₃	2	N(CH ₃) ₂	321-323	C ₁₉ H ₂₂ N ₄ O.2HCl	-1.02
14	NH ₂	5-OCH ₃	2	N(CH ₃) ₂	>360	C ₁₉ H ₂₂ N ₄ O ₂ .2HCl	-1.06
15	NH ₂	5-Cl	2	N(CH ₃) ₂	311-312	C ₁₈ H ₁₉ ClN ₄ O.2HCl	
16	NH ₂	6-NO ₂	2	N(CH ₃) ₂	>360	C ₁₈ H ₁₉ N ₅ O ₃ .2HCl	
17	NH ₂	6-NH ₂	2	N(CH ₃) ₂	>360	C ₁₈ H ₂₁ N ₅ O.2HCl	
18	NH ₂	6-CH ₃	2	N(CH ₃) ₂	326-328	C ₁₉ H ₂₂ N ₄ O.2HCl	-0.82
19	NH ₂	6-OCH ₃	2	N(CH ₃) ₂	256-258	C ₁₉ H ₂₂ N ₄ O.2HCl	
20	NH ₂	7-NO ₂	2	N(CH ₃) ₂	316-318	C ₁₈ H ₁₉ N ₅ O ₃ .2HCl	-1.29
21	NH ₂	7-NH ₂	2	N(CH ₃) ₂	324-326	C ₁₈ H ₂₁ N ₅ O.3HCl	-1.64
22	NH ₂	7-CH ₃	2	N(CH ₃) ₂	316-319	C ₁₉ H ₂₂ N ₄ O.2HCl	
23	NH ₂	7-OCH ₃	2	N(CH ₃) ₂	290-292	C ₁₉ H ₂₂ N ₄ O ₂ .2HCl $\frac{1}{2}$ H ₂ O	
24	NH ₂	7-Cl	2	N(CH ₃) ₂	310-311	C ₁₈ H ₁₉ ClN ₄ O.2HCl	

BEST AVAILABLE COPY

- 16 -

TABLE II

Elemental analyses for the compounds of Table I

0098098

No	Formula	Found				Calculated			
		C	H	N	Cl	C	H	N	Cl
1	$C_{18}H_{19}N_3O \cdot 2HCl$	59.1	5.9	11.7	19.0	59.0	5.8	11.5	19.4
2	$C_{19}H_{21}N_3O \cdot 2HCl$	60.3	6.0	11.0	18.5	60.0	6.1	11.1	18.6
3	$C_{19}H_{22}N_4O \cdot 2HCl$	57.6	6.4	14.1	18.5	57.7	6.1	14.2	17.9
4	$C_{18}H_{20}N_4O \cdot 2HCl \cdot \frac{1}{2} H_2O$	56.0	6.2	15.1	18.4	56.0	5.9	14.5	18.4
5	$C_{19}H_{22}N_4O \cdot 2HCl$	58.1	6.3	14.6		57.7	6.1	14.2	
6	$C_{18}H_{20}N_4O_2 \cdot 2HCl$	54.2	5.7	14.0	17.6	54.4	5.6	14.1	17.8
7	$C_{20}H_{24}N_4O \cdot 2HCl$	59.1	6.5	13.6	17.1	58.7	6.4	13.7	17.3
8	$C_{16}H_{16}N_4O \cdot 2HCl$	54.5	5.1	16.0	19.8	54.4	5.1	15.9	20.1
9	$C_{18}H_{19}N_5O_3 \cdot 2HCl$	50.5	4.8	16.4	16.8	50.7	5.0	16.4	16.6
10	$C_{18}H_{21}N_5O \cdot 2HCl$	52.0	5.9	16.7		52.2	6.1	16.9	
11	$C_{18}H_{19}N_5O_3 \cdot 2HCl$	50.5	5.2	16.5	16.6	50.7	5.0	16.4	16.6
12	$C_{18}H_{21}N_5O \cdot 2HCl \cdot H_2O$	52.6	6.0	17.1		52.2	6.1	16.9	
13	$C_{19}H_{22}N_4O \cdot 2HCl$	58.2	6.2	14.2	17.9	57.7	6.1	14.2	17.9
14	$C_{19}H_{22}N_4O_2 \cdot 2HCl$	55.5	6.4	13.2	16.5	55.5	5.9	13.6	17.2
15	$C_{18}H_{19}ClN_4O \cdot 2HCl$	52.0	5.0	13.4	25.2	52.0	5.1	13.5	25.6
16	$C_{18}H_{19}N_5O_3 \cdot 2HCl$	50.6	5.2	16.4	16.5	50.7	5.0	16.4	16.6
17	$C_{18}H_{21}N_5O \cdot 2HCl$	54.6	5.6	17.8	17.8	54.5	5.9	17.7	17.9
18	$C_{19}H_{22}N_4O \cdot 2HCl$	57.8	6.1	14.3	17.9	57.7	6.1	14.2	17.9
19	$C_{19}H_{22}N_4O_2 \cdot 2HCl$	55.1	6.1	13.8	17.7	55.4	5.9	13.6	17.3
20	$C_{18}H_{19}N_5O_3 \cdot 2HCl$	51.0	4.9	16.3	16.1	50.7	5.0	16.4	16.6
21	$C_{18}H_{21}N_5O \cdot 3HCl$	49.9	5.6	16.2	24.6	50.0	5.5	16.4	24.2
22	$C_{19}H_{22}N_4O \cdot 2HCl$	57.4	5.9	14.0	17.8	57.7	6.1	14.2	17.9
23	$C_{19}H_{22}N_4O \cdot 2HCl \cdot \frac{1}{2} H_2O$	54.4	6.2	13.1	16.5	54.3	6.0	13.3	16.9
24	$C_{18}H_{19}ClN_4O \cdot 2HCl$	52.5	4.7	13.5	25.4	52.0	5.1	13.5	25.6

The following Examples illustrate the preparation of compounds of the general formula (I):

EXAMPLE A: Preparation of compound 4 of Table I by the method of Scheme I

5 N-(2-Carboxyphenyl)anthranilic Acid (IV, R₂=H)

A mixture of 2-chlorobenzoic acid (100g), anthranilic acid (90g), anhydrous powdered K₂CO₃ (135g), Cu/CuO (2g) and 2-ethoxyethanol (200ml) was heated with swirling on the steam bath until gas evolution ceased and then stirred at 145° for a further 2 1/2 hours.

10 The thick reaction mixture was diluted with water, acidified (HCl) and then the crude product was collected and washed well with hot water. This was dissolved in hot dilute aqueous Na₂CO₃ treated liberally with charcoal-celite and filtered through a celite pad. The hot filtrate was diluted with half the volume of
15 EtOH and then slowly acidified (HCl). The pale yellow product which separated was collected when still warm, washed well with hot water, benzene and dried, providing material (84% yield) of sufficient purity for use in the next step (lit, m.p. 295° dec.).

20 9(10H)Acridone-4-carboxylic Acid (V, R₂=H)

A mixture of the preceding diphenylamine diacid (80g) and conc. H₂SO₄ (250ml) was heated at 100°C for 4 hours, then cooled, poured into ice-water and the precipitated solid collected and washed well with water. This was dissolved in dilute aqueous

25 NaOH and following filtration was diluted with an equal volume of EtOH and then acidified with glacial acetic acid (this left any sulfonated impurities in solution). The acridone acid which slowly crystallized from the hot solution was collected after thorough cooling, washed with EtOH, water, EtOH again and dried
30 providing pure material in 83% yield m.p. 342-343° dec.

9-Chloroacridine-4-carbonyl chloride (II, R₂=H, Z=Cl; R₆=Cl)

A suspension of the preceding acridone acid (20g) in SOCl₂ (60ml) containing DMF (2 drops) was heated gently under reflux with

stirring until homogeneous and then for a further 45 min. The solution was evaporated to dryness in vacuo, below 40°C, and residual traces of SOCl₂ were removed by addition of dry benzene and complete re-evaporation of all solvents to give the crude
5 product as a yellow powder.

N-[2-(dimethylamino)ethyl] 9-chloroacridine-4-carboxamide (II,
R₂=H, Z=Cl, R₆=NH(CH₂)₂ N(CH₃)₂.

The above carbonyl chloride was cooled to -5° and to this was added in one portion an ice-cold solution of
10 N,N-dimethylethylenediamine (36.5ml) in dry dichloromethane (200ml). After stirring at 30°C until homogeneous the reaction solution was left for a further 15min and then shaken with dilute aq. Na₂CO₃. The organic layer was washed with dilute aq. Na₂CO₃ (2x), aq. NaCl solution and then dried (Na₂SO₄). Evaporation of
15 the solvent left an oil which slowly solidified. This was extracted with hot dry benzene-petroleum ether (1:5), treated with charcoal-celite and filtered quickly through a hot celite pad. Crystalline material rapidly separated and addition of further petroleum ether completed precipitation of the product.
20 The yellow solid was collected, washed with petroleum ether and dried providing material 19.5g (71% yield) indicated by TLC to contain only trace quantities of the corresponding acridone and this product was stored over KOH and used without further purification.

25 Compound 4 of Table 1

The above compound (II; R₂=H, Z=Cl, R₆=NH(CH₂)₂N(CH₃)₂ (4.0g) was dissolved in dry phenol (12.8g) and heated slowly to 50°C, to provide a solution of the phenoxy compound (II; R₂=H, Z=OC₆H₅), R₆=NH(CH₂)₂ NH(CH₃)₂) in excess phenol. A stream of dry ammonia
30 was passed into the solution while the temperature was raised from 50°C to 115°C. Addition of ammonia was continued for 15 min, after which the mixture was cooled and diluted with excess 40% aqueous NaOH. Prolonged cooling gave a solid that was crystallized from aqueous EtOH and then EtOAc. The resulting

pure base was converted to the dihydrochloride salt by dissolving in MeOH, treating with 12N HCl (2.2 equivalents) and precipitating with EtOAc. Crystallization from MeOH/EtOAc gave hygroscopic yellow prisms of the pure dihydrochloride of compound 4, m.p. 304-305°C (72% yield).

Minor modifications of the procedure of Example A, employing appropriately substituted 2-chlorobenzoic acids and/or appropriate amine components, were used to prepare compounds 3, 5, 7, 8, 9-15 and 18 of Table I. Separation of isomers after ring closure was sometimes required.

EXAMPLE B: Preparation of compound 1 of Table 1 by the method of Scheme I.

4-Carboxyacridine (VI; Z=H, R₂=H)

4-Carboxyacridanone (V; R₂=H) (5g) and NaOH (1g; 1.1 equivalent) were dissolved in water (100ml). Al foil (3g; amalgamated by dipping each piece into a solution of 15g of mercuric chloride in 100ml of water for 5 min immediately before use) was added in pieces to the stirred, boiling solution of carboxyacridanone over 30min. After a further 30min reflux, the hot solution was filtered and acidified with HCl. FeCl₃ (12g) was added, and the mixture was heated until clear (an initial heavy precipitate redissolves) and for a further 10mins. The mixture was basified with 2N NaOH, filtered from Fe(OH)₃, and the pH adjusted to 5, when a precipitate formed. This was collected, washed with water and extracted with boiling EtOH (400ml). The filtrate was concentrated to 30ml and cooled well, yielding 4-carboxyacridine (2.2g, 47%), m.p. 202-204°C.

Compound 1 of Table I

4-Carboxyacridine (1.1g, 5.6mM) was refluxed in SOCl₂ (10ml) and a drop of DMF for 1h, and the volatiles were evaporated. Dry benzene (20ml) was added and evaporated to remove residual traces of SOCl₂, and the resulting solid was dissolved in dry DMF (20ml) containing N,N-dimethylethylenediamine (1.25g, 3 equivalents).

The mixture was kept at 20°C for 2h and the volatiles were evaporated at 40°C. The resulting gum was extracted with boiling diisopropyl ether, and this solution was concentrated and diluted with petroleum ether to give the free base as yellow needles
5 (0.85g, 61%). The free base was dissolved in MeOH and dry HCl gas added to pH2. Dilution with EtOAc gave the dihydrochloride as yellow crystals (87%), m.p. 195-197°C.

EXAMPLE C: The preparation of compounds 20 and 21 of Table I by the methods of Scheme I and Scheme II.

10 N-(2-methoxycarbonylphenyl)-5-nitroanthranilic acid (VIII, R₂ = 7-NO₂, R₇ = CH₃)

A mixture of 2-chloro-5-nitrobenzoic acid (7g, 35mM), methyl anthranilate (6.3g, 45mM), and cupric acetate (6.3g, 35mM) in bis-isopropylethylamine (10ml) and N-methylpyrrolidone (5ml) was
15 stirred and heated at 150° for 2h under N₂. The cooled solution was diluted with water and acidified with 2N HCl. The gummy precipitate was collected by decantation and triturated with a small amount of cold MeOH to give a yellow solid. This was collected and washed with cold MeOH to give the desired diphenylamine ester
20 (VIII; R₂ = 7-NO₂, R₇ = CH₃) (2.6g, 24%). Crystallization from EtoAc gave yellow needles, m.p. 228-229°C.

Methyl 7-nitro-acridanone-4-carboxylate (IX, R₂ = 7-NO₂, R₇ = CH₃)

The above diphenylamine ester (2.0g) was heated at 100°C for 1h
25 with polyphosphate ester (10g). The cooled mixture was diluted with water and basified with Na₂CO₃ to give the acridone ester, which was collected and crystallized from EtOH as yellow needles, m.p. 310-312°C (1.7g, 91% yield).

7-Nitro-4-carboxyacridanone (V, R₂=7-NO₂)

30 The above acridone ester (2.0g) was heated in 92% H₂SO₄ (50ml) for 7h at 100°C. The cooled mixture was poured into water and the precipitate collected and extracted with aqueous Na₂CO₃. The

extract was filtered and acidified with 2N HCl to provide pure product (1.72g, 90%), which was recrystallized from DMF as a yellow powder, m.p. about 375°C.

7-Nitro-4-carboxyacridanone (V; R₂ = 7-NO₂) (by direct nitration).

- 5 A stirred solution of 4-carboxyacridanone (10.0g) in c.H₂SO₄ (50ml) was treated portionwise at below 5°C with powdered KNO₃ (4.6g), then stirred for 30min at 20°C and poured into ice water. The precipitate was collected, washed, dried and crystallized from DMF/MeOH and then DMF to give pure product of m.p. about
10 375°C (65% yield).

The product was identical (assessed by TLC) to the compound obtained above by polyphosphate ester ring closure of N-(2-methoxycarboxylphenyl)-5-nitroanthranilic acid and subsequent acid hydrolysis of the methyl ester function.

15 Compound 20 of Table I

- The above 7-nitro-4-carboxyacridanone was converted to 7-nitro-9-chloroacridine-4-carbonyl chloride (II, Z=Cl, R₂ = 7-NO₂, R₃ =Cl), and treated with N,N-dimethylethylenediamine followed by dry ammonia in phenol by the methods outlined in
20 Example A above to give N-(2-dimethylaminoethyl)-9-amino-7-nitroacridine-4-carboxamide dihydrochloride (compound 20 of Table I), m.p. 316-318°C.

Compound 21 of Table I

- The above nitro compound was reduced using Fe powder and HCl in
25 65% aqueous EtOH. Basification with 2N NaOH gave the crude product, which was converted to the trinydrochloride with 12N HCl in MeOH/EtOAc. Two recrystallizations from MeOH/EtOAc gave pure product, m.p. 324-326°C.

EXAMPLE D: Preparation of compound 6 of Table I by the method of Scheme I

6-p-Nitrophenyl acridanone-4-carboxylate. (VII; R₂=H)

Pure, finely-powered acridanone-4-carboxylic acid (19.8g, 83mM) and p-nitrophenol (22.2g, 160mM) were suspended in pyridine (200ml). The mixture was stirred vigorously at 60°C while PCl₃ (4.4ml; 53mM) was added dropwise. The mixture was immediately heated to 100°C until homogeneous. On cooling product separated, and after 1h the reaction was cooled well and the precipitate collected and washed well with acetone. Recrystallization from DMF gave pure compound (74% yield), m.p. 280-281°C.

p-Nitrophenyl 9-chloroacridine-4-carboxylate (II; Z=Cl, R₂=H, R₆=OC₆H₄NO₂).

The above compound (2.02g, 5.6mM) was refluxed gently in SOCl₂ (6ml) and a drop of DMF for 1h. The volatiles were evaporated, dry benzene was added, and the volatiles evaporated again to remove all traces of HCl and SOCl₂. The residue was dissolved in CH₂Cl₂, cooled to 0°C, and icecold 10% KHCO₃ (20ml) was added. The organic layer was separated, dried, and concentrated to small volume to provide the product as yellow needles (80% yield), m.p. 194-196°C.

Compound 6 of Table I

4-Nitrophenyl 9-chloroacridine-4-carboxylate (0.01 M) was added in one portion to an ice-cooled stirred solution of 2-(2-aminoethylamino)-ethanol (0.012 M) and triethylamine (0.011 M) in anhydrous dichloromethane (20ml). The mixture was stirred until homogeneous, and then for a further 10 min. Dry phenol (11g) was then added to the solution and anhydrous ammonia was passed in while the temperature was raised to 115°C. After contact with ammonia at this temperature for a further 10 min the mixture was cooled and excess 5N aqueous NaOH added. The resulting solid was dissolved in 1N aqueous HCl and this solution was slowly neutralised with 1N aqueous NH₄OH, precipitating a quantity of material that was removed by filtration and

discarded. Treatment of the filtrate with excess aqueous NaOH gave crude material which was recycled through the above purification process. The resulting free base was crystallized from MeOH-H₂O. Crystallisation of the dihydrochloride salt from MeOH-EtOAc then provided pure product, m.p. 292-293°C dec.

EXAMPLE E: Preparation of compound 2 of Table I by the method of Scheme V.

2-(N-2-Methylcarbonylphenyl)aminobenzoic acid (XII; R₂=H, R₈=OH)

A mixture of 2-chloroacetophenone (20g, 0.18mol), anthranilic acid (37g, 0.27mol), dry K₂CO₃ (37g, 0.27mol), Cu powder (0.1g), and CuCl (0.1g) suspended in 50ml dimethoxyethane was stirred under reflux for 20h and cooled. The mixture was extracted with dilute aqueous NaOH, the solution was clarified with charcoal-celite, filtered, and acidified to give 7.4g (16%) of 2-(N-2-methylcarbonylphenyl)aminobenzoic acid, (XII; R₂=H, R₈=OH) m.p. 280-283° decomp. (EtOH).

N-(2-Dimethylaminoethyl)-2-(N-2-methylcarbonylphenyl) aminobenzamide (XII; R₂=H, R₈=NH(CH₂)₂ N(CH₃)₂)

A solution of the above keto-acid in dry DMF (20ml) was treated with 2.3g (1.2 equivalents) of diethyl phosphorocyanidate (DEPC), and an excess of N,N-dimethylethylenediamine (2g) was added dropwise. After being warmed on a waterbath for 30min the reaction mixture was basified with K₂CO₃ (aq) and the solvent was removed under vacuum. The residue was extracted with ethyl acetate, and after being washed (H₂O, brine) and dried (Na₂SO₄) the solvent was removed to give crude

N-(2-dimethylaminoethyl)-2-(N-2-methylcarbonylphenyl) aminobenzamide, (XII; R₂=H, R₈=NH(CH₂)₂ N(CH₃)₂) as an oil.

Compound 2 of Table I

The above oily product was dissolved in 25ml of a mixture of 100 parts HOAc and 5 parts H₂SO₄. After heating under reflux for 1h the HOAc was removed under vacuum and the residue was dissolved in water. The aqueous fraction was extracted with CH₂Cl₂.

basified with aqueous NaOH and extracted with EtOAc. The EtOAc layer was washed with water and saturated brine, dried and evaporated to provide

- 5 N-(2-dimethylaminoethyl)-9-methylacridine-4-carboxamide (2.1g, 60%) as an oil. This was dissolved in MeOH-EtOAc and treated with dry HCl gas to provide compound 2 of Table I as a crystalline dihydrochloride salt, m.p. 178-180°C (recrystallized from EtOH).

10 Alternative Preparation of Compound 2 of Table I by the method of Scheme IV

Ethyl 9-methylacridine-4-carboxylate (XIII; R₂=H, R₈=OEt).

- To a solution of 1.2g (4.7mmol) of 2-(N-2-methylcarbonylphenyl) aminobenzoic acid in 5ml dry DMF was added 1.15g DEPC (1.5 equivalents), ethanol (1ml) and Et₃N (1.4g, 3 equivalents) and
15 the mixture was heated on a water bath for 1h. Since the reaction was incomplete a further equivalent of each reagent was added and the mixture was heated for a further 1h. The solvent was removed under vacuum, and the residue was basified with KHC0₃(aq) and extracted into EtOAc. After washing and drying
20 (Na₂SO₄) the solvent was removed to give the crude ethyl ester (XII; R₂=H, R₈=OEt) which was dissolved in 15ml of a mixture of HOAc (100 parts) and H₂SO₄ (5 parts). After heating under reflux for 1h the HOAc was removed under vacuum and the residue was basified with dil KHC0₃ solution, and extracted with EtOAc. The
25 organic layer was washed with dilute aqueous methanesulphonic acid and discarded, and after being basified with dil. KHC0₃ solution the aqueous layer was extracted with EtOAc to give 0.31g of ethyl 9-methylacridine-4-carboxylate (XIII; R₂=H, R₈=OEt) (23%) as an oil.

30 N-(2-Dimethylaminoethyl)-9-methylacridine-4-carboxamide.
(Compound 2 of Table I).

The ethyl ester from above (1mmol) was treated with refluxing 1N NaOH in 20ml 60% aqueous ethanol for 1h; the solution was neutralized by the dropwise addition of conc. HCl and the solvent was

- removed under vacuum. The crude acid (XIII; $R_2=H$, $R_8=OH$) was then treated with refluxing $SOCl_2$ for 30min, and after removal of the solvent the product acid chloride (XIII; $R_2=H$, $R_8=Cl$) was dissolved in dry CH_2Cl_2 . The solution was cooled in ice and an excess of N,N-dimethylethylenediamine was then added slowly. The solution was then washed well with water to remove excess amine and after being dried (Na_2SO_4) the solvent was removed to give N-(2-dimethylaminoethyl)-9-methylacridine-4-carboxamide (0.24g, 78%) as an oil. This oil was converted to the crystalline hydrochloride salt (compound 2 of Table I), m.p. 178-180° decomp (EtOH).

EXAMPLE F: Preparation of Compound 16 of Table I by the method of Scheme II

5-Nitrodiphenyliodonium-2-carboxylate

- 2-Iodo-4-nitrobenzoic acid (17.5g, 0.06mol) (W.C. Agosta, Tet. Lett., 1965, 2681) was dissolved in 60ml conc H_2SO_4 and the solution was cooled to 0°C. Potassium persulphate (31.2g, 0.116mol) was added in portions with stirring over 40 min, and after a further 60min at <10°C 55ml of benzene was added. The stirred viscous mixture was then allowed to warm slowly to room temperature and left overnight. The reaction mixture was then poured onto ice, and the white precipitate was filtered off and suspended in a stirred solution of 200ml 5N NaOH. After being filtered off and washed several times with water the solid was dried by azeotroping with benzene. Yield 20.1g, 91%. The product was insoluble in all of the normal solvent systems but has some solubility in DMF.

N-[2-(Methoxycarbonyl)phenyl]-4-nitro-anthranilic acid (VIII)

- Crude 5-Nitrodiphenyliodonium-2-carboxylate (42g, 0.114mol) was suspended in 500ml of DMF containing 34g methyl anthranilate and 1.0g $Cu(OAc)_2$, and the mixture was heated on a waterbath for 2 days when all of the solid had dissolved. The dark red-brown solution was diluted firstly with 50ml of conc NH_3 , and then with 2L of water and the oily insolubles were removed by washing twice

with dichloromethane. Acidification with dilute HCl then gave the nitro ester acid, 31.3g, 86%, which was recrystallized from EtOAc as red needles, m.p. 243-245°C.

Methyl 3-nitro-9(10H)-acridanone-5-carboxylate (IX)

- 5 The above half ester (1.0g, 3.2mM) was heated with polyphosphate ester at 100°C for 1h. The cooled product was diluted with water and basified to pH 9. The insoluble product was collected and crystallized from ethanol as yellow prisms: m.p. 252-253°C.

3-Nitro-5-carboxy-9(10H)-acridanone (V)

- 10 The above acridone ester (1.5g, 5.0 mM) was heated in sulphuric acid (20ml, 92% v/v) for 7h at 100°C. The cooled mixture was diluted with water, and the product collected and washed. Trituration with 2N aq Na₂CO₃ followed by removal of insoluble products and acidification of the filtrate gave the acid (1.29g, 15 90% yield). A sample was crystallized from a large volume of EtOH, : m.p. 375°C. Attempted ester hydrolysis under basic conditions gave impure, deeply coloured products.

Compound 16 of Table I

- The above 4-carboxyacridanone was converted, via the
20 9-chloro-4-carbonyl chloride to compound 16 of Table I, using the methods given in Example A.

EXAMPLE G: Preparation of compound 23 of Table I by the method of Scheme III

2-(4-Methoxyphenylamino)-1,3-benzenedicarboxylic acid (XI)

- 25 A mixture of 10g 2-iodoisophthalic acid (34 mmol), 8.3g p-anisidine (70mmol), 0.5g CuCl, 0.5g Cu(OAc)₂, 10ml N-ethylmorpholine and 25ml DMF was heated with stirring at 125°C for 2h under nitrogen and cooled. The reaction mixture was then diluted with 100ml of dil HCl and extracted with EtOAc. The
30 organic layer was extracted with dil NaOH solution and discarded. Acidification of the aqueous layer with dil HCl gave a precipitate of the methoxy-diacid, 8.64g, 88% m.p. 226-228°C (EtOAc).

2-Methoxy-acridanone-5-carboxylic acid (V)

The above diacid (4.3g, 30mM) was treated with polyphosphoric acid (30g) for 4h at 130°C. The cooled melt was dissolved in water and the pH was adjusted to 7.5 with aqueous NaOH to precipitate the crude acridone acid.

Compound 23 of Table I

The above crude acridanone acid was treated as outlined in Example A to provide compound 23 of Table I.

The procedure of Example G was also used with appropriate choice of starting materials to prepare compounds 19, 22 and 24 of Table I.

The compounds of general formula (I), and particularly the examples listed in Tables I and II, have antitumour activity in both in vivo and in vitro test systems, as shown by the data of Table III. This Table gives biological data for compounds 1-24, whose physical data has been given in Tables I and II. The abbreviations given in Table III are:-

- | | | |
|----|-----------------------|---|
| 10 | P388 <u>in vivo</u> | - Tumour P388 cells were obtained as frozen stocks from Mason Research Inc., U.S.A. and passaged intraperitoneally according to standard methods (<u>Cancer Chemother. Rep.</u> 3, Part 3, page 9, 1972) in DBA-2 mice of either sex. Groups of six F1 hybrid mice (DBA-2 male x C57 B1 female, g weight 20 ± 1 g) were injected intraperitoneally with 10^6 cells on day 0. |
| 15 | O.D. | - optimal drug dose, in milligrams per kilogram, administered as a solution in 0.1 ml of 30% v/v ethyl alcohol in water on days 1, 5 and 9 after tumour inoculation. The drug is administered as a soluble acid addition salt. |
| 20 | ILS | - percentage increase in life span of treated animals over that of groups of control animals injected with tumour alone. The average survival of control mice was 11 days. Values of ILS greater than 20% are considered statistically significant. |
| 25 | | |
| 30 | L1210 <u>in vitro</u> | - The culture methods used are described in detail elsewhere (B.C. Baguley and R. Nash, <u>Europ.J.Cancer</u> , 17, 671-679 (1981). Acceptable reproducibility of data depends critically upon the maintenance of optimal culture conditions. L1210 cells were ini- |

tially obtained from Dr I. Wodinsky, Arthur D. Little Inc., Boston, U.S.A., under the auspices of the National Cancer Institute.

ID₅₀

5

- the nanomolar concentration of drug which, when added to cultures of murine L1210 leukaemic cells over a period of 70 hours, reduces the resultant counted number of leukaemia cells by 50% (B.C. Baguley and R. Nash, Europ.J.Cancer, 17, 671-679 (1981)). Values below 1000nM are considered significant.

10

The compounds of general formula (I) also show broad-spectrum antibacterial activity. Specifically, compound 4 is active against the bacteria *Aerobacter aerogenes*, *Alcaligenes viscolactis*, *Escherichia coli*, *Bacillus subtilis*, *Sarcina lutea*, *Micrococcus lysodeikticus*, *Neisseria catarrhalis*, *Staphylococcus aureus*, *Xanthomonas phaseoli* and *Streptococcus faecalis*.

15

Biological data for the compounds of Table I 0098098

No.	L1210 in vitro	P388 in vivo		
		ID ₅₀	OD	ILS Active
1	105	66	91	Y
2	66	45	14	N
3	15	5.9	53	Y
4	15	4.5	98	Y
5	157	20	0	N
6	77	20	80	Y
7	5.5	5.9	70	Y
8	414	20	71	Y
9	319	8.9	25	Y
10	162	30	28	Y
11	1.3	0.8	39	Y
12	18	8.9	58	Y
13	0.33	2.6	107	Y
14	4.3	3.9	81	Y
15	2.9	2.6	81	Y
16	0.05	2.6	23	Y
17	35	8.9	58	Y
18	55	2.6	20	Y
19	151	8.9	17	Y
20	104	20	34	Y
21	48	20	80	Y
22	605	13.3	0	N
23	518	13.3	4	N
24	722	13.3	8	N

The active compounds may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or they may be enclosed in hard or soft shell gelatin capsules, or they may be compressed into tablets, or they may be incorporated directly with the food of the diet. For oral therapeutic administration, the active compounds may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers and the like. Generally, such compositions and preparations should contain at least 0.1% of active compound. The percentage in the compositions and preparations may, of course be varied and may conveniently be from about 2% to about 60% of the weight of the unit. The amount of active compound in such therapeutically use-

ful compositions is such that a suitable dosage will be obtained. Preferred compositions or preparations according to the present invention are prepared so that an oral dosage unit form contains from about 5 to about 200 milligrams of active compound.

- 5 The tablets, troches, pills, capsules and the like may also contain, for example, one or more of the following:
a binder such as gum tragacanth, acacia, corn
starch or gelatin; excipients such as dicalcium phosphate; a
disintegrating agent such as corn starch, potato starch, alginic
10 acid and the like; a lubricant such as magnesium stearate; and a
sweetening agent such a sucrose, lactose or saccharin may be
added or a flavouring agent such as peppermint, oil of win-
tergreen or cherry flavouring. When the dosage unit form is a
capsule, it may contain, e.g., in addition to materials of the above
15 type, a liquid carrier. Various other materials may be present
as coatings or to otherwise modify the physical form of the
dosage unit. For instance, tablets, pills or capsules may be
coated with shellac, sugar or both. A syrup or elixir may con-
tain the active compound, sucrose as a sweetening agent, methyl
20 and propylparabens as preservatives, a dye and flavouring such as
cherry or orange flavour. Of course, any material used in pre-
paring any dosage unit form should be pharmaceutically pure and
substantially non-toxic in the amounts employed. In addition,
the active compounds may be incorporated into sustained-release
25 preparations and formulations.

The active compounds may also be administered parenterally or intraperitoneally. Solutions of the active compound as a free base or pharmaceutically acceptable salt can be prepared in water suitably mixed with a surfactant such a hydroxypropylcellulose.

- 30 Dispersions can also be prepared e.g. in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.



The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be
5 fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol
10 (for example, glycerol, propylene glycol, liquid polyethylene glycol, and the like), suitable mixtures thereof and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use
15 of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium
20 chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminium monostearate and gelatin.

Sterile injectable solutions may be prepared by incorporating the
25 active compound in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic
30 dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze-drying technique which yield a powder of the active ingredient plus any additional

desired ingredient from a previously sterile-filtered solution thereof.

As used herein, "pharmaceutically acceptable carrier" includes any and all suitable solvents, dispersion media, coatings, antibacterial
5 and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active
10 ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.


It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to
15 physically discrete units suitable as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the novel dosage unit
20 forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active material and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active material for the treatment of disease in living subjects having a diseased
25 condition in which bodily health is impaired as herein disclosed.

The principal active ingredient may be compounded for convenient and effective administration in effective amounts with a suitable pharmaceutically acceptable carrier in dosage unit form as
30 hereinbefore disclosed. A unit dosage form can, for example, contain the principal active compound in amounts ranging from about 0.1 to about 400 mg. with from about one to about 30 mg being preferred. Expressed in proportions, the active compound

is generally present in from about 0.1 to about 400 mg/ml of carrier. In the case of compositions containing supplementary active ingredients, the dosages are determined by reference to the usual dose and manner of administration of the said ingredients.

5

1. A compound represented by the general formula (I),




Y represents $C(NH)NH_2$, $NHC(NH)NH_2$, or NR_4R_5 , where each of R_4 and R_5 , which may be the same or different, represents H or lower alkyl unsubstituted or substituted by one or more of the same or different substituents selected from hydroxyl and amino functions; and


or an acid addition salt thereof.

2. A compound according to Claim 1 where R_1 represents NH_2 , R_2 represents up to two of 1-, 5-, 6-, 7-, and 8- NO_2 , 5- and 6- CH_3 , and 5-Cl, Y represents $NHC(NH)NH_2$, $N(CH_3)_2$ or $NHCH_2CH_2OH$ and x is 2.
- 5 3. A compound according to Claim 1 where R_1 represents H, R_2 represents up to two of 1-, 5-, 6-, 7-, and 8- NO_2 , 5- and 6- CH_3 , and 5-Cl, Y represents $NHC(NH)NH_2$, $N(CH_3)_2$ or $NHCH_2CH_2OH$ and x is 2.
- 10 4. A compound according to Claim 1 in which R_1 and R_2 represent H, Y represents $N(CH_3)_2$ and x is 2.

5. A compound according to Claim 1 in which R_1 represents NH_2 , R_2 represents H, Y represents $N(CH_3)_2$ and x is 2.
6. A compound according to Claim 1 in which R_1 represents NH_2 , R_2 represents 6- NO_2 , Y represents $N(CH_3)_2$ and x is 2.
- 5 7. A compound according to Claim 1 in which R_1 represents NH_2 , R_2 represents 5- CH_3 , Y represents $N(CH_3)_2$ and x is 2.
8. A process for the preparation of a compound represented by the general formula (I), as defined in Claim 1, or an acid addition salt thereof, which comprises coupling a substituted acridine of the general formula (II),
- 10


II

where R_2 represents groups as defined in Claim 1, Z represents H, CH_3 or a suitable leaving group and R_6 represents Cl, Br or OC_6H_4 -p- NO_2 , with a primary alkyl amine of the general formula (III),

III


- 15 where x and Y are defined as in Claim 1, and, when Z is a leaving group, converting the resultant coupled product to a compound of general formula (I) where R_1 represents NHR_3 and R_3 is as defined in Claim 1, and, if desired, converting a compound of formula (I) into an acid addition salt thereof.

9. A process according to Claim 8 wherein Z in formula (II) represents a leaving group selected from methoxy, phenoxy, alkylthio, and halogen.
10. A process according to Claim 8 wherein Z in formula (II) represents chloro.
11. A process according to any one of Claims 8 to 10 wherein R₆ in formula (II) represents Cl or OC₆H₄-p-NO₂.
12. A process according to any one of claims 8 to 11 wherein the coupling reaction of compound (II) with compound (III) is performed in an anhydrous solvent selected from chloroform, dimethylsulphoxide, N-methylpyrrolidone, dichloromethane, and dimethylformamide, buffered with a tertiary amine.
13. A process according to any one of Claims 8 to 12 wherein Z in formula (II) represents chloro and the resultant coupled product is treated with anhydrous ammonia or amine of the formula R₃NH₂ in phenol or cresol to provide a compound of the formula (I) where R₁ represents NHR₃ and R₃ is defined in Claim 1.
14. A process for the preparation of a compound represented by the general formula (I), as defined in Claim 1, in which R₁ represents CH₃, or an acid addition salt thereof, which comprises cyclodehydrating a compound of the general formula (XII')



where x and y are defined as in Claim 1, and R₂'

- and R_2 " represent hydrogen or together represent up to 2 of the groups CH_3 , OCH_3 , halogen, CF_3 , NO_2 , NH_2 , $NHCOCH_3$ and $NHCOOCH_3$ placed at positions corresponding to positions 1 to 3 and 5 to 8
- 5 in the formula (I), and, if desired, converting a compound of formula (I) into an acid addition salt thereof.
15. A compound represented by the general formula (I), as defined in Claim 1, or an acid addition salt thereof, whenever prepared by the process according to any one of Claims 8 to 13.
- 10
16. A compound of the general formula (II)




where R_2 represents H or up to two of the groups CH_3 , OCH_3 , halogen, CF_3 , NO_2 , NH_2 , $NHCOCH_3$, and $NHCOOCH_3$ placed at positions 1-3 and 5-8;

Z represents H, CH_3 or a leaving group selected from methoxy, phenoxy, alkylthio, and halogen; and

R_6 represents Cl, Br or $OC_6H_4-p-NO_2$, or an acid addition salt thereof.

17. A compound of the general formula (IX)




IX

- where R₂ represents H or up to two of the groups CH₃, OCH₃, halogen, CF₃, NO₂, NH₂, NHCOCH₃, and NHCOOCH₃ placed at positions 1-3 and 5-8; and
- 5 R₇ represents a lower alkyl group, or an acid addition salt thereof.
18. A pharmaceutical preparation having antitumour activity which comprises at least one compound of the general formula (I) defined in Claim 1, or a pharmaceutically acceptable acid addition salt thereof, in admixture or conjunction with one or more pharmaceutically acceptable carriers or diluents.
- 10
19. A pharmaceutical preparation having antitumour activity which comprises a compound according to any one of Claims 2 to 7 and 15, in admixture or conjunction with a pharmaceutically acceptable acid addition salt thereof, and one or more pharmaceutically acceptable carriers or diluents.
- 15
20. A compound of the general formula (I) as defined in Claim 1, or a pharmaceutically acceptable acid addition salt thereof, for use in the treatment of tumours.
- 20

21. A compound according to any one of Claims 2 to 7 and 15, or a pharmaceutically acceptable acid addition salt thereof, for use in the treatment of tumours.

CLAIMS FOR AUSTRIA:

1. A process for the preparation of a compound represented by the general formula (I),




5 where R_1 represents H, CH_3 or NHR_3 , where R_3 represents H, $COCH_3$, SO_2CH_3 , $COPh$, SO_2Ph or lower alkyl unsubstituted or substituted by one or more of the same or different substituents selected from hydroxyl and amino functions;

10 R_2 represents H or up to two of the groups CH_3 , OCH_3 , halogen, CF_3 , NO_2 , NH_2 , $NHCOCH_3$ and $NHCOOCH_3$ placed at positions 1-3 and 5-8;


15 Y represents $C(NH)NH_2$, $NHC(NH)NH_2$, or NR_4R_5 , where each of R_4 and R_5 , which may be the same or different, represents H or lower alkyl unsubstituted or substituted by one or more of the same or different substituents selected from hydroxyl and amino functions; and

x is from 2 to 6,

20 or an acid addition salt thereof, which process comprises coupling a substituted acridine of the general formula (II),

II


where R_2 represents groups as defined above, Z represents H , CH_3 or a suitable leaving group and R_6 represents Cl , Br or $\text{OC}_6\text{H}_4\text{-p-NO}_2$, with a primary alkyl amine of the general formula (III),

III

- 5 where x and Y are defined as above, and, when Z is a leaving group, converting the resultant coupled product to a compound of general formula (I) where R_1 represents NHR_3 and R_3 is as defined in Claim 1, and, if desired, converting a compound of formula (I) into an acid addition salt thereof.
- 10 2. A process according to Claim 1 wherein Z in formula (II) represents a leaving group selected from methoxy, phenoxy, alkylthio, and halogen.
3. A process according to Claim 1 wherein Z in formula (II) represents chloro.
- 15 4. A process according to any one of Claims 1 to 3 wherein R_6 in formula (II) represents Cl or $\text{OC}_6\text{H}_4\text{-p-NO}_2$.
5. A process according to any one of claims 1 to 4 wherein the coupling reaction of compound (II) with compound (III) is performed in an anhydrous solvent selected from chloroform, dimethylsulphoxide, N-methylpyrrolidone, dichloromethane, and
- 20 dimethylformamide, buffered with a tertiary amine.
6. A process according to any one of Claims 1 to 5 wherein Z in formula (II) represents chloro and the resultant coupled product is treated with anhydrous ammonia or amine of the

formula R_3NH_2 in phenol or cresol to provide a compound of the formula (I) where R_1 represents NHR_3 and R_3 is defined in Claim 1.

7. A process for the preparation of a compound represented by the general formula (I), as defined in Claim 1, in which R_1 represents CH_3 , or an acid addition salt thereof, which comprises cyclodehydrating a compound of the general formula (XII')




- where x and Y are defined as in Claim 1, and R_2' and R_2'' represent hydrogen or together represent up to 2 of the groups CH_3 , OCH_3 , halogen, CF_3 , NO_2 , NH_2 , $NHCOCH_3$ and $NHCOOCH_3$ placed at positions corresponding to positions 1 to 3 and 5 to 8 in the formula (I), and, if desired, converting a compound of formula (I) into an acid addition salt thereof.

8. A process according to any one of Claims 1 to 7 in which R_1 represents NH_2 , R_2 represents up to two of 1-, 5-, 6-, 7-, and 8- NO_2 , 5- and 6- CH_3 , and 5- Cl , Y represents $NHC(NH)NH_2$, $N(CH_3)_2$ or $NHCH_2CH_2OH$ and x is 2.

9. A process according to any one of Claims 1 to 7 in which R_1 represents H , R_2 represents up to two of 1-, 5-, 6-, 7-, and 8- NO_2 , 5- and 6- CH_3 , and 5- Cl , Y represents $NHC(NH)NH_2$, $N(CH_3)_2$ or $NHCH_2CH_2OH$ and x is 2.

10. A process according to any one of Claims 1 to 7 in which R_1 and R_2 represent H, Y represents $N(CH_3)_2$ and x is 2.
11. A process according to any one of Claims 1 to 7 in which R_1 represents NH_2 , R_2 represents H, Y represents $N(CH_3)_2$ and x is 2.
12. A process according to any one of Claims 1 to 7 in which R_1 represents NH_2 , R_2 represents 6- NO_2 , Y represents $N(CH_3)_2$ and x is 2.
13. A process according to any one of Claims 1 to 7 in which R_1 represents NH_2 , R_2 represents 5- CH_3 , Y represents $N(CH_3)_2$ and x is 2.
14. A process for the preparation of a compound represented by the general formula (II)




II

15 where R_2 is defined as in Claim 1;

Z represents H, CH_3 , Cl, or Br; and


R_6 represents Cl, Br, or $OC_6H_4-p-NO_2$,

or an acid addition salt thereof, which process comprises reacting a compound of the general formula (VI)




VI

where R_2 is defined as above; and Z represents H, or CH_3 , with a halogenating agent or with tris(4-nitrophenyl)phosphite or reacting a compound of the general formula (V)



V

- 5 where R_2 is defined as above, with a halogenating agent or with tris(4-nitrophenyl)phosphite and then with a halogenating agent, and, if desired, converting a compound of formula (II) into an acid addition salt thereof.
- 10 15. A process for the preparation of a compound represented by the general formula (IX)




IX

where R_2 is defined as in Claim 1; and

R_7 represents a lower alkyl group,

or an acid addition salt thereof, which process comprises ring closing a compound of the general formula (VIII)



5 in which R_2 and R_7 are defined as above,
and if desired, converting a compound of formula
(IX) into an acid addition salt thereof.

- 10 16. A process for the preparation of a pharmaceutical preparation having antitumour activity which comprises bringing into admixture or conjunction at least one compound of the general formula (I) defined in Claim 1, or a pharmaceutically acceptable acid addition salt thereof, and one or more pharmaceutically acceptable carriers or diluents.
- 15 17. Use of a compound of the general formula (I) as defined in Claim 1, or a pharmaceutically acceptable acid addition salt thereof, in the treatment of tumours.

19



Europäisches Patentamt
European Patent Office
Office européen des brevets

11 Publication number:

0 098 098
A3

12

EUROPEAN PATENT APPLICATION

21 Application number: 83303610.6

22 Date of filing: 22.06.83

51 Int. Cl.⁴: C 07 D 219/04
C 07 D 219/10, C 07 D 219/06
A 61 K 31/435

30 Priority: 25.06.82 NZ 201084

43 Date of publication of application:
11.01.84 Bulletin 84/2

88 Date of deferred publication of search report: 02.05.85

84 Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

71 Applicant: DEVELOPMENT FINANCE CORPORATION OF
NEW ZEALAND
Development Finance Centre Corner Grey and
Featherston Streets
Wellington(NZ)

72 Inventor: Atwell, Graham John
37 Hawkins Street Meadowbank
Auckland 5(NZ)

72 Inventor: Baguley, Bruce Charles
36a Olsen Avenue Hillsborough
Auckland 4(NZ)


72 Inventor: Denny, William Alexander
165 Gossamer Drive Pakuranga
Auckland(NZ)

72 Inventor: Rewcastle, Gordon William
24 Rothery Road Manurewa
Auckland(NZ)

74 Representative: Senior, Janet et al,
Abel & Imray Northumberland House 303-306 High
Holborn
London WC1V 7LH(GB)

54 Acridinecarboxamide compounds.

57 The novel class of 4-carboxamidoacridines of the pre-
sent invention represented by the general formula (I),



where

R₁ represents H, CH₃ or NHR₃, where R₃ is H, COCH₃,
SO₂CH₃, CPh, SO₂Ph or lower alkyl optionally substituted
with hydroxyl and/or amino functions;

R₂ represents H or up to two of the groups CH₃, OCH₃,
halogen, CF₃, NO₂, NH₂, NHCOCH₃, and NHCOOCH₃ placed
at positions 1-3 and 5-8;

Y represents C(NH)NH₂, NHC(NH)NH₂, or NR₄R₅, where
each of R₄ and R₅ is H or lower alkyl optionally substituted
with hydroxyl and/or amino functions; and

x is from 2 to 6,

and the acid addition salts thereof, possess antibacterial and
antitumour properties.

EP 0 098 098 A3

Croydon Printing Company Ltd



European Patent
Office

EUROPEAN SEARCH REPORT

0098098

Application number

EP 83 30 3610

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 3)
D, A P	EP-A-0 073 155 (DEVELOPMENT FINANCE CORPORATION OF NEW ZEALAND) * Claims 1,21,22 *	1, 17, 18	C 07 D 219/04 C 07 D 219/10 C 07 D 219/06 A 61 K 31/435
A	EP-A-0 039 224 (DEVELOPMENT FINANCE CORPORATION OF NEW ZEALAND) * Claims 1,6 *	1, 18	
A	Chemical Abstracts vol. 78, no. 4, 29. Januar 1973 Columbus, Ohio, USA K. BURDESKA et al. "Synthesis of 1-substituted 4-nitroacridones", page 86, column 1, abstract no. 17591b & Helvetica Chimica Acta vol. 55, no. 6, 1972, page s 1948-1958 in combination with Chemical Substance Index 1972-1976, page 775 CS, column 1, lines 5-7	17	TECHNICAL FIELDS SEARCHED (Int. Cl. 3) A 61 K 31/435 C 07 D 219/04 C 07 D 219/06 C 07 D 219/10
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 30-10-1984	Examiner KNAACK M
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO Form 1503 03 82